

Review article

A review of phytochemical of biological studies of *Cycas* species growing in Middle East

Ahmed Ismail ¹

¹ Pharmacognosy Department, Faculty of Pharmacy, Fayoum University, Fayoum, Egypt; ais03@fayoum.edu.eg; https://orcid.org/0000-0003-0266-8760

RECEIVED 16 December 2024
 REVISED 17 January 2025
 ACCEPTED 27 January 2025
 ONLINE 11 April 2025

CITATION

Ismail, A (2025). A review of phytochemical of biological studies of *Cycas* species growing in Middle East. *Spectrum Science Journal*, 2(1): 1-16
 DOI: 10.21608/sasj.2025.345074.1001

CORRESPONDENCE

Ahmed Ismail
 Email: ais03@fayoum.edu.eg

The Scientific Association for Studies and Applied Research (SASAR)
 https://sasj.journals.ekb.eg



COPYRIGHT

©2024 by the authors. Submitted for possible open access publication under the terms and conditions of the Creative Commons Attribution (CC BY) license.

Spectrum Science Journal
 PRINT ISSN: 3062-5068
 ONLINE ISSN: 3062-5076

Abstract:

The genus *Cycas*, commonly known as cycads, represents a fascinating group of ancient gymnosperms with a lengthy evolutionary history that dates back to the Mesozoic era. The taxonomy, morphology, ecology, and conservation status of the genus *Cycas* are all covered in detail in this abstract. A review of phytochemical constituents and biological activities of *Cycas* species has been presented. Notably, while there are approximately 580 species of this genus distributed worldwide, only nine species grow in the Middle East, and just three of these have been studied phytochemically. *Cycas* species are of particular interest due to their unique evolutionary position, ecological roles, and potential medicinal properties. This review provides a comprehensive overview of the genus *Cycas*, integrating biological activity, chemical diversity, traditional knowledge, and economic and ecological significance. By synthesizing research across these fields, we seek to create a holistic view of the species, highlighting its historical, current, and potential future significance, relevance in the realms of ecology, ethnobotany, and medicine. Hence, this review highlights the phytochemical studies of *Cycas* species, underscoring their potential as a source of bioactive molecules with applications in medicine and beyond.

Keywords: *Cycas*, species, economic, phytochemical, traditional, pharmacological.



INTRODUCTION

For thousands of years, herbal remedies have played a significant role in the treatment and management of a wide range of diseases and improvement of the quality of human life. A large numbers of drugs, commonly used worldwide, are nowadays derived throughout scientifically based investigation of traditionally used remedies, especially medicinal plants. This trend facilitates medication with natural products rather than synthetic analogues, which could be harmful. The potential of higher plants as a source for new drugs is still largely unexplored. A relatively small number has been phytochemically investigated and the fraction submitted to biological or

pharmacological screening is even smaller (1). According to taxonomy, there are more than a hundred species of *Cycas*, which are found in tropical and subtropical climates. They exhibit a great diversity in terms of leaf form, reproductive structures, and preferred habitats. *Cycas* species names are related to palms; the term "palm" in Greek means kykas, is where they got their name. The aerial trunk of *Cycas* plants is columnar, and their leaf crowns are pinnately complex. The stem is usually unbranched. a spiral-shaped crown of leaves at the top of the underground tuberous stems of immature plants. Cones are spirally aggregated reproductive organs made of highly modified leaves called sporophylls, which are produced by cyclops. Numerous sporangia, or pollen capsules, are carried by each male sporophyll and are often found on the underside. Although every female sporophyll is home to ovules, usually two (2). *Cycas* is known to be the only genus of family Cycadaceae and represented in Egypt (3), by nine species; *Cycas armstrongii* Miq., *Cycas revoluta* Thunb., *Cycas circinalis* L., *Cycas litoralis* K.D. Hill, *Cycas thouarsii* R.Br., *Cycas media* R.Br, *Cycas tansachana* K.D. Hill, *Cycas rumphii* Miq. and *Cycas pectinata* Griff. (4). In China and Japan, some available pharmaceuticals used for treatment of cancer and hepatitis contain *Cycas revoluta* Thunb. and *C. circinalis* L. plants (5), They were also directed for anticancer, antimicrobial, antimalarial and antileishmanial activities.

Taxonomy of *Cycas*

Kingdom	Plantae
Subkingdom	Tracheobionta
Division	Cycadophyta
Class	Cycadopsida
Order	Cycadales
Family	Cycadaceae
Genus	<i>Cycas</i>
Species	<i>Cycas armstrongii</i> Miq. <i>Cycas circinalis</i> L. <i>Cycas revoluta</i> Thunb. <i>Cycas litoralis</i> <u>K.D. Hill</u> <i>Cycas media</i> R.Br <i>Cycas pectinata</i> Griff <i>Cycas rumphii</i> Miq. <i>Cycas tansachana</i> <u>K.D. Hill</u> <i>Cycas thouarsii</i> R.Br.

Morphology

Cycas is the only genus currently recognized in the cycad family Cycadaceae. The most well-known species is *Cycas revoluta*, which is commonly grown under the names "sago palm, Japanese cycad, Sotetsu, king sago palm" despite not actually being a palm. The generic name comes from Greek kykas and means "palm tree". The species of this genus are concentrated around the equatorial regions. The plants are dioecious, and the family Cycadaceae is unique among the cycads in not forming seed cones on female plants, but rather a group of leaf-like structures each with seeds on the lower margins, and pollen

cones on male individuals. *Cycas* plants are dioecious palm like trees with aerial or subterranean, cylindrical stems.

The caudex is cylindrical, surrounded by the persistent petiole base. Most species form distinct branched or unbranched trunks but in some species the main trunk can be subterranean with the leaf crown appearing to arise directly from the ground. Spirally arranged, pinnate or very rarely bipinnate, lower leaflets are frequently reduced to spines, and leaves are produced in seasonal growth flushes intermingled with cataphylls. Horizontal ptyxis circinate, longitudinal ptyxis erect or infrequently reflexed. Most species' stomata are restricted to the abaxial surface of the leaflets, which have a single, thick midrib and no lateral veins. Each ptyxis is involute. Pubescent leaves with simple or branching transparent hairs, at least when they are young. Each microsporophyll has a simple, sterile apex that frequently develops into an inverted spine. Microsporophylls are aggregated into determinate male cones. On its abaxial surfaces, every microsporophyll has a large number of microsporangia, or pollen sacs. Microsporangia with slits for opening. Megasporophylls are spirally arranged in an indeterminate terminal rosette, with the central axis continuing vegetative growth, and are loosely or tightly imbricate. Two to many (rarely one) ovules that are obliquely pointed outward (ascending), marginally implanted on the stipe. In the distal zone beyond the ovule-bearing stipe, megasporophyll apically dilated into a pinnatifid, pectinate, toothed, or whole lamina. The seeds are subglobular to ellipsoidal, and beneath the inner woody sclerotesta, there may or may not be spongy tissue. The outer sarcotesta is fleshy and can be yellow, orange, or brown. Endosperm haploid, generated from the female gametophyte. Embryo is straight, features a long, spirally twisted suspensor and two cotyledons that are typically connected at the tips. platyspermic seeds with cryptocotylar germination. Photographs of *Cycas* species growing in middle east are shown in Figure.1.

Ecological Significance of genus *Cycas*:

Cycads have an ancient fossil record, they are considered as living representatives of an ancient plant lineage and their interactions with animals (13). They play an important role in nitrogen fixation in soils and enhancing their fertility by the association with cyanobacteria, its commonly known as endosymbiosis (14).

Economic and Ethnobotanical significances of genus *Cycas*:

Indigenous tribes around the world have used the numerous species of the genus *Cycas* for a wide range of customary uses. Many elements of human life have benefited from the use of these ancient plants, also known as cycads, from food and medicine to cultural and ceremonial rituals. These are a few noteworthy traditional use for various *Cycas* species.

□ Nutrition:

Sago palms, or *Cycas revoluta*, are among the most well-known *Cycas* species because of their starchy pith, which is taken out of the center of the trunk. For many indigenous populations in Southeast Asia and the Pacific Islands, processed sago has long been a staple diet. It is a substantial source of carbohydrates that is processed, extracted, and added to a number of traditional meals. (15), Cakes made in Sri Lanka are made from the starch extracted from its seeds (16).

□ Traditional Medicine:

Cycas rumphii pollen grains are used to reduce discomfort and ease pain. *Cycas circinalis* seed paste is applied topically to heal wounds, boils, swellings, and skin conditions and is used to heal ulcers. For the purpose of preventing blood vomiting, young *Cycas revoluta* leaves are juiced, while blood vomiting, stomach issues, and many skin problems are treated using extract from its young plants. *Cycas revoluta* tincture made from its seeds is applied to certain conditions like as headache, nausea, and sore throat. A purgative is made by decocting seeds. Ulcerated sores and swollen glands are treated with *Cycas circinalis* terminal buds. Hair roots are treated with the stem of *Cycas pectinata* (16, 17).

. Photographs of *Cycas* species growing in middle east are shown in Figure.1.



Figure 1. Photograph for *Cycas* species in middle east (1): *Cycas armstrongii* Miq., (2): *Cycas circinalis* L., (3): *Cycas revoluta* Thunb., (4): *Cycas littoralis* K.D. Hill ., (5): *Cycas media* R.Br, (6): *Cycas pectinata* Griff ., (7): *Cycas rumphii* Miq., (8): *Cycas tansachana* K.D. Hill ., (9): *Cycas thouarsii* R.Br. (6-12)

□ Cultural and Ritual Importance:

Different *Cycas* Species: For indigenous groups, a variety of *Cycas* species have ceremonial and cultural significance (18) . Certain plants are included into ceremonies, rituals, and customs in some

societies because they are revered. There can be symbolic significance to the existence of particular *Cycas* species in particular places.

□ Constructions and fibers source:

Native American tribes have utilized some *Cycas* species, especially those with dense, sturdy wood, for building projects. These species provide wood that can be used to make tools, create traditional shelters, or even build ceremonial buildings. Traditional textiles have been woven using fibres that taken from the leaves or trunks of some *Cycas* species. These fabrics can be used to make baskets, apparel, and other objects of functional and cultural value. its leaves are utilised to make baskets, hats and mats (16).

□ Insects Repellent:

Many *Cycas* species like *Cycas sphaerica* are used, as insect repellent properties have long been associated with the fronds of this plant. Frequently burned, these fronds produce compounds that repel pests and insects, offering a useful defense against annoyances (19).

□ Ornamental purposes: Cycads often referred to as cycads, are well-liked options for ornamental landscaping because of their unique look, hardiness, and flexibility. Many species in the genus *Cycas* are grown for their beautiful appearance, which gives gardens, parks, and public areas a hint of ancient charm (20). Cycads are valuable decorative plants in areas with limited water supplies or sporadic water restrictions since they are typically resistant to drought conditions. They can endure in dry conditions because of their ability to hold water in their pithy stems. In addition, they can be found growing in a variety of soil types, such as clay, sandy, and loamy soils. Their versatility in landscaping projects is increased by this adaptability, which qualifies them for a range of garden designs and soil types; they are prized for their lengthy lifespans and for giving landscapes a timeless beauty as ornamental plants. These plants' slow growth guarantees that they keep their visual appeal for a long time, which helps to create sturdy and long-lasting garden elements in Urban areas (21). Sago palm or *Cycas revoluta*, one of the most extensively grown varieties, is the main applications of *Cycas* as an ornamental plant and is valued for its eye-catching and perfectly symmetrical feather-like fronds. Because of its glossy, deep green foliage, which create an exotic, tropical atmosphere, it's a popular choice for landscaping (22).

Chemical composition of genus *Cycas*:

The diverse range of chemicals and metabolic processes found in the chemistry of *Cycas* species, popularly referred to as cycads, have developed over millions of years. Since they are ancient gymnosperms, cycads provide a unique window into the chemical adaptations that have enabled these plants to flourish in a variety of global environments, predating even flowering plants. A fascinating story of natural chemicals, metabolic pathways, and ecological interactions is revealed by the chemistry of *Cycas* species, from their historic uses in indigenous societies to their contemporary applications in pharmacology and ecology. *Cycas* species exhibit a wide range of secondary metabolites and a vast repertory of phytochemical constituents. These comprise phenolic chemicals, alkaloids, flavonoids, and terpenoids, each of which adds to the distinct chemical signature of various *Cycas* species. The investigation of these phytochemicals offers important new information for a number of scientific fields in addition to providing evidence for the evolutionary tactics employed by cycads. Chemical investigation of the constituents of genus *Cycas* resulted in isolation and identification of including different classes as flavonoids, phenolic acids, catechins and sterols. All constituents are listed in Table (1), Figure (2).

Table 1: List of reported compounds from different *Cycas* species

<i>C. armstrongii</i> Miq.	Phytochemistry	N-(3'-one-5'-methyl)-hexyl-alanine, β -sitosterol, Stigmasterol, Naringenin, Kaempferol aglycone, Dihydroamentoflavone, Dihydrobilobetin, 2,3-Dihydrohinokiflavone, Amentoflavone, β -sitosterol glucoside, Isoginkgetin, Prunin, Naringin, Vanillic acid, P-Coumaric acid, 4-Acetoxybenzoic acid butyl ester, Isopimara-7,15-dien-19-ol, Isopimara-7,15-diene, Phthalic acid dibutyl ester, Pregnane-3,12,14,20-tetrol, Pinoresinol, Glanduloidin C, NeoCycasin A, NeoCycasin J, Lanosterol, Salvileucolide methyl ester, β -carotene.(4). Vitexin, Vitexin-2''-rhamnoside, Caffeine (6). Catechin, Epicatechin, Epigallo-catechin 3-gallate (14)
	Biology	Antimicrobial, antileshmanial, antitrypanosomal, antioxidant and an ameliorative role on γ radiation toxicity in brain and pancreas (4)
<i>C. circinalis</i> L.		Glucosyloxyazoxymethane (Cycasin) (23), Methylazoxymethanol (24), Caffeic acid, p-hydroxybenzoic acid, Vanillic acid, p-coumaric acid, Protocatechuic acid and Ferulic acid (25)
	Phytochemistry	N-(3'-one-5'-methyl)-hexylalanine and Leucine betaine (26) Naringenin, Dihydroamentoflavone, Dihydrobilobetin, Dihydroisoginkgetin, Tetrahydrobilobetin, Tetrahydroisoginkgetin, Amentoflavone, Bilobetin, Isoginkgetin, Dihydrohinokiflavone, Vitexin-glucoside, Vicenin-2, Catechin, Epicatechin, Gallocatechin, Epigallocatechin, Methyl glucose, Loliolide, Dihydrodehydrodiconiferyl alcohol, β -Sitosterol glucoside (27), α -amino-b-methylaminopropionic acid (28)
	Biology	Isopimara-7,15-dien-19-ol, Isopimara-7,15-diene, 9,12-Dihydroxy-15-nonadecenoic acid, Pinoresinol, NeoCycasin A, NeoCycasin J, Lanosterol, Salvileucolide methyl ester, β -carotene, Kaempferol-3-O-Rhamnosyl-(1 \rightarrow 2)-galactoside (4) Antibacterial (29), immunomodulatory activity (30) Antimicrobial, antileshmanial, antitrypanosomal, antioxidant and an ameliorative role on γ radiation toxicity in brain and pancreas (4)
<i>C. revoluta</i> Thunb.	Phytochemistry	Sotetsuflavone (31), Caffeic acid, p-hydroxybenzoic acid, Vanillic acid, p-coumaric acid, Protocatechuic acid and Ferulic acid (25). Sugars; Arabinose, Galactose, Fucose, Rhamnose, Methylrhamnose and Mannose (32). Cycasinidene, Cycasthiamide (33) Cycasin and Macrozamin (34). Minerals; Ca, Mg, Fe, Zn (35) Polyphenols; Naringenin, Prunin, Dihydroamentoflavone, Dihydroamentoflavone monoglucoside,

		Dihydroamentoflavone diglucoside, Tetrahydroamentoflavone, Amentoflavone, Podocarpusflavone A, Hinokiflavone, Dihydrohinokiflavone, Dihydroisocryptomerin, Tetrahydrohinokiflavone, Vitexin rhamnoside, Vomifoliol, Lariciresinol, Isolariciresinol, Protocatechuic acid (27). 5,6,7,8,3',4'-hexamethoxyflavone , 5,6,7,8,4-pentamethoxyflavone, 7,3'-dihydroisoflavones (36) 4-Acetoxybenzoic acid butyl ester, Isopimara-7,15-diene, Isopimara-7,15-dien-19-ol, 5,8-Dihydroxy-9,12-octadecadienoic acid, 5-hydroxy-6,7,8-trimethoxy-flavanone, NeoCycasin A, NeoCycasin J, Lanosterol, Salvileucolide methyl ester, β -carotene, kaempferol 3-O-rutinoside (4)
	Biology	Colon cancer protective activity (37), antimicrobial (38), cytotoxic and antioxidant (39). Antileishmanial, antitrypanosomal, antioxidant and an ameliorative role on γ radiation toxicity in brain and pancreas (4)
<i>C. litoralis</i> K.D. Hill	Phytochemistry Biology	No reports
<i>C. media</i> R.Br	Phytochemistry Biology	a-amino-b-methylaminopropionic acid (40), Sequoyitol (41) and Macrozamin (42) Anti-Biofilm and antibacterial Activities (7)
<i>C. pectinata</i> Griff.	Phytochemistry Biology	Amentoflavone and 2,3-dihydroamentoflavone (8) Anti-diabetic (8)
<i>C. rumphii</i> Miq.	Phytochemistry Biology	Amentoflavone (43), Cycasindene (10) Caffeic acid , p-hydroxybenzoic acid, Vanillic acid, P-coumaric acid, Protocatechuic acid and Ferulic acid (25) Antibacterial (11)
<i>C. tansachana</i> K.D. Hill	Phytochemistry Biology	No reports
<i>C. thouarsii</i> R.Br.	Phytochemistry Biology	Shikimic acid, Quinic acid, Chlorogenic acid, Trigonelline, Ferulic Acid, Rosmarinic acid, Quercetin 3-O-glucuronide, Homoorientin, Hesperetin-7-O-neohesperidoside, Vitexin-2''-O-rhamnoside, Delphinidin 3-rutinoside, Vitexin, Luteolin-7-O-glucoside, Kaempferol-7-neohesperidoside, Luteolin, Isorhamnetin., Prunin, Apigenin-7-O-glucoside , Peonidin-3-glucoside, Quercetin-3-Glucuronide, Naringenin, Apigenin, Genistein (12). Antibacterial activity and wound healing (12)




Figure 2: Structures of reported compounds from different *Cycas* species







Figure 2. (Cont.)




Cycasindene




Cycathioamide




Leucine betaine




N-(3'-one-5'-methyl)-hexyl-alanine




α -amino- β -methylaminopropionic acid (L-BMAA)




R= H MAM
R= B-D-glucose Cycasin




Methylaminoalanine




β -sitosterol



β -sitosterol glucoside




Lanosterol




Stigmasterol


Figure 2. (Cont.)




Pregnane-3,12,14,20-tetrol




Salvileucolide methyl ester




Epigallo-catechin 3-gallate




Caffeine




Catechin



Epicatechin




Gallocatechin




Epigallocatechin


Figure 2. (Cont.)




Methyl glucose




Sequoyitol




Loliolide




Dihydrodehydrodiconiferyl alcohol




Vomifoliol



Lariciresinol




Isolariciresinol




Dihydroaemontoflavone monoglucoside
Dihydroaemontoflavone diglucoside

R= H
R= β -D-glucopyranosyl



β -carotene



Macrozamin

Figure 2. (Cont.)













Figure 2. (Cont.)

Conclusion

Cycas plants are abundantly found in tropical and subtropical regions of the world. The species are used for their pharmacologically established properties, including those against cancer, tumors, broad-spectrum microbes, viruses, and fungi, have all been described and demonstrated. Several additional verification of pharmacological efficacy are necessary, even though the physiological illnesses share common symptoms. Out of nine species of this genus distributed in Middle East, just three species have been phytochemically investigated. Phytochemical work on this genus seems to be by no means exhaustive and there still remains a vital scope for study of active molecules. Biflavonoids are widely distributed throughout this genus which suggests that these compounds may be chemotaxonomic markers of the genus *Cycas*.

Finally, the genus *Cycas* is revealed as a botanical treasure trove that is not limited by time. The review's compilation of collective knowledge encourages more multidisciplinary investigation and emphasizes the value of comprehensive strategies for comprehending, protecting, and maximizing *Cycas*' potential. This review highlights *Cycas* species as a potential source of bioactive molecules. As we traverse the delicate interplay between ecological resilience, cultural richness, and chemical variety within this species, we are reminded of the urgency to maintain and appreciate the priceless contributions of *Cycas* to the natural and cultural legacy of our world.

Methodology

Limited phytochemical studies on *Cycas* Middle Eastern species made a challenge for collecting enriched data. A deep search in many websites as Google Scholar, Research gate with aid of a database library engine of The Egyptian Knowledge Bank; EKB.

CRedit authorship contribution statement

AI: Conceptualization, Writing – original draft, data curation, Investigation, Validation.

Declaration of Competing Interest: The author declares that he has no competing interests.

Funding: No funding resources.

References

1. Minhas FA, Rehaman H-u, Yasin A, Awan ZI, Ahmed N. Antimicrobial activities of the leaves and roots of *Elaeagnus umbellata* Thunb. *African Journal of Biotechnology*. 2013;12(48):6754-60.
2. El-Seadawy HM, Abo El-Seoud KA, El-Aasr M, Ragab AE. Cycadaceae: An Important Source for Biflavonoids and Various Pharmacological Effects of Different *Cycas* Species. *Journal of Advanced Medical and Pharmaceutical Research*. 2023.
3. Moawad A, Hetta M, Zjawiony JK, Jacob MR, Hifnawy M, Marais JP, et al. Phytochemical investigation of *Cycas circinalis* and *Cycas revoluta* leaflets: moderately active antibacterial biflavonoids. *Planta medica*. 2010;76(08):796-802.
4. Ismail A, Hassan HM, Moawad AS, Abdel Fattah SM, Sherif NH, Abdelmohsen UR, et al. Chemical composition and therapeutic potential of three *Cycas* species in brain damage and pancreatitis provoked by γ -radiation exposure in rats. *Journal of Radiation Research and Applied Sciences*. 2020;13(1):200-14.
5. Wang K. New formulations of traditional Chinese medicine for treating AIDS. . Faming Zhuanli Shenqing Gongkai Shuomingshu, CN 101085325 A 20071212. 2007.
6. Ismail A. Phytochemical and biological studies of some *Cycas* species cultivated in Egypt. PhD Thesis, Beni-Suef Faculty of Pharmacy. 2020.
7. Attallah NG, Al-Fakhrany OM, Elekhawy E, Hussein IA, Shaldam MA, Altwaijry N, et al. Anti-biofilm and antibacterial activities of *Cycas media* R. Br secondary metabolites: In silico, in vitro, and in vivo approaches. *Antibiotics*. 2022;11(8):993.

8. Laishram S, Sheikh Y, Moirangthem D, Deb L, Pal B, Talukdar N, et al. Anti-diabetic molecules from *Cycas pectinata* Griff. traditionally used by the Maiba-Maibi. *Phytomedicine*. 2015;22(1):23-6.
9. *Cycas pectinata*" Text available under a CC-BY-SA Creative Commons Attribution License. 2005 [Available from: http://www.llifile.com/Encyclopedia/PALMS_AND_CYCADS/Family/Cycadaceae/28863/Cycas_pectinata.
10. Pan M, Mabry TJ, Cao P, Moini M. Identification of nonprotein amino acids from cycad seeds as N-ethoxycarbonyl ethyl ester derivatives by positive chemical-ionization gas chromatography-mass spectrometry. *Journal of Chromatography A*. 1997;787(1-2):288-94.
11. Khan AV, Ahmed QU, Khan AA, Shukla I. Antibacterial activity of *Cycas rumphii* Miq. leaves extracts against some tropical human pathogenic bacteria. *Research Journal of Microbiology*. 2011;6(10):761.
12. Binsuwaidan R, Elekhawey E, Elseady WS, Keshk WA, Shoeib NA, Attallah NG, et al. Antibacterial activity and wound healing potential of *Cycas thouarsii* R. Br n-butanol fraction in diabetic rats supported with phytochemical profiling. *Biomedicine & Pharmacotherapy*. 2022;155:113763.
13. Hall J. The ecology of cycads: Living representatives of an ancient plant lineage and their interactions with animals. 2011.
14. Ismail A, Radwan MM, Hassan HM, Moawad AS, Hetta MH. Isolation of catechins from *Cycas armstrongii* Miq. of an Egyptian origin. *Journal of Applied Pharmaceutical Science*. 2021;11(1):158-62.
15. Li K, Zhang T, Zhao W, Ren H, Hong S, Ge Y, et al. Characterization of starch extracted from seeds of *Cycas revoluta*. *Frontiers in Nutrition*. 2023;10:1159554.
16. Trisha. *Cycas*: Occurrence, Morphological Features and Economic Importance [Available from: <https://www.biologydiscussion.com/articles/Cycas-occurrence-morphological-features-and-economic-importance/5720>.
17. Darade MS. Gymnospermic Medicines Used In Disease Treatment. *World Journal of Pharmaceutical and Life Sciences*. 2022;8(8):238-43.
18. Bonta M, Bamigboye S. Use of cycads as ritual and recreational narcotics. *South African Journal of Botany*. 2018;115:280-1.
19. Raju AS, Jonathan KH. Anemophily, accidental cantharophily, seed dispersal and seedling ecology of *Cycas sphaerica* Roxb.(Cycadaceae), a data-deficient red-listed species of northern Eastern Ghats. *Current Science*. 2010:1105-11.
20. Goel AK, Khuraijam J. Cycads: an overview. *Plant Biology and Biotechnology: Volume I: Plant Diversity, Organization, Function and Improvement*. 2015:349-60.
21. Khuraijam J, Roy R. Indian *Cycas* under severe threat. *Current Science*. 2016;111(11):1756-8.
22. Forrester MB, Layton GM, Varney SM. *Cycas revoluta* (sago cycad) exposures reported to Texas poison centers. *The American Journal of Emergency Medicine*. 2020;38(8):1611-5.
23. Riggs N. Glucosyloxyazoxymethane, a constituent of the seeds of *Cycas-circinalis* L. SOC CHEMICAL INDUSTRY 14 BELGRAVE SQUARE, LONDON SW1X 8PS, ENGLAND; 1956. p. 926-.
24. Kobayashi A, Matsumoto H. Studies on methylazoxymethanol, the aglycone of *Cycasin*: Isolation, biological, and chemical properties. *Archives of biochemistry and biophysics*. 1965;110(2):373-80.
25. Wallace JW. A survey for benzoic and cinnamic acids of the Cycadaceae. *American Journal of Botany*. 1972;59(1):1-4.
26. Li C-J, Brownson DM, Mabry TJ, Perera C, Bell EA. Nonprotein amino acids from seeds of *Cycas circinalis* and *Phaseolus vulgaris*. *Phytochemistry*. 1996;42(2):443-5.
27. Moawad AS. Comparative Phytochemical and Biological Study of Certain Plants of Family Cycadaceae Cultivated in Egypt. PhD Thesis, Beni-Suef Faculty of Pharmacy. 2011.
28. Nunn PB. 50 years of research on α -amino- β -methylaminopropionic acid (β -methylaminoalanine). *Phytochemistry*. 2017;144:271-81.
29. Raslin A, Rajkumar J. Antimicrobial activity of a polyherbal formulation Ambrex. *International Journal of Materials and Product Technology*. 2017;55(1-3):286-95.

30. Shilpashree V, Dang R, Das K. Evaluation of phytochemical investigation and immunomodulatory activity of four different plant species of vidari by carbon clearance test on wister rats. *ANNALS OF PHYTOMEDICINE-AN INTERNATIONAL JOURNAL*. 2015;4(1):94-8.
31. Kawano N, Yamada M. The Structure of Sotetsuflavone. *Journal of the American Chemical Society*. 1960;82(6):1505-6.
32. De Luca P, Moretti A, Sabato S, Gigliano GS. A comparative study of cycad mucilages. *Phytochemistry*. 1982;21(7):1609-11.
33. Pan M, Mabry TJ, Beale JM, Mamiya BM. Nonprotein amino acids from *Cycas revoluta*. *Phytochemistry*. 1997;45(3):517-9.
34. Yagi F. Azoxyglycoside content and β -glycosidase activities in leaves of various cycads. *Phytochemistry*. 2004;65(24):3243-7.
35. Qiu-Li Y. Determination of Trace Elements in *Cycas Revoluta* by Flame Atomic Absorption Spectrometry [J]. *Chinese Journal of Spectroscopy Laboratory*. 2007;5.
36. Xu Y, Dai Q-C, Luan Z-Z, Xiang Z, Wu J, Fan Y-L, editors. *The Isolation and Identification Compounds from Cycas Revoluta*. *Medicine and Biopharmaceutical: Proceedings of the 2015 International Conference*; 2016: World Scientific.
37. Bera S, Das B, De A, Samanta A. Preventive effect of *Cycas revoluta* in 1, 2-Dimethylhydrazine-induced colon cancer in wistar rat model. *Asian J Pharm Clin Res*. 2018;11(5):120-3.
38. Lal K, Ahmed N, Mathur A. Study of the Antimicrobial Profile and Phytochemical Composition of Solvent Extracts of Leaves and Female Cones of *Cycas revoluta*. *Int J Curr Microbiol App Sci*. 2017;6(4):2514-22.
39. Negm WA, Ibrahim AE-RS, El-Seoud KA, Attia GI, Ragab AE. A New Cytotoxic and Antioxidant Amentoflavone Monoglucoside from *Cycas revoluta* Thunb Growing in Egypt. *Journal of Pharmaceutical Sciences and Research*. 2016;8(5):343.
40. Duncan MW, Kopin IJ, Crowley JS, Jones SM, Markey SP. Quantification of the putative neurotoxin 2-amino-3-(methylamino) propanoic acid (BMAA) in cycadales: analysis of the seeds of some members of the family Cycadaceae. *Journal of analytical toxicology*. 1989;13(3):169-75.
41. Riggs N. Virtual coupling in the cyclic six-spin system of sequoyitol. *Australian Journal of Chemistry*. 1964;17(5):603-6.
42. Riggs N. The occurrence of macrozamin in the seeds of cycads. *Australian Journal of Chemistry*. 1954;7(1):123-.
43. Uddin Q, Malik A, Azam S, Hadi N, Azmi A, Parveen N, et al. The biflavonoid, amentoflavone degrades DNA in the presence of copper ions. *Toxicology in vitro*. 2004;18(4):435-40.



Open Access Journal

© 2025 by the authors. Disclaimer/Publisher's Note: The content in all publications reflects the views, opinions, and data of the respective individual author(s) and contributor(s), and not those of the scientific association for studies and applied research (SASAR) or the editor(s). SASAR and/or the editor(s) explicitly state that they are not liable for any harm to individuals or property arising from the ideas, methods, instructions, or products mentioned in the content.